

Oblique Lumbar Interbody Fusion (OLIF)

WHAT DOES SURGERY ENTAIL?

This is a surgery where we make an incision on your side of your abdomen. After splitting the muscle, we gently move things around to access the spine from the front or the side. With a special retractor and instruments, we can safely protect your organs and blood vessels. We then remove the disk between the vertebrae in its entirety and replace it with a cage or spacer which increases the space between the bones. This helps to remove pressure off the nerves that have been compressed due to a degenerative disk. The cage has special bone graft material which, over the course of about a year, helps the bones heal through the cage completing the fusion. We then place screws and rods through your back through small poke holes to act as an internal brace to hold everything together while your bones heal through the cage. Sometimes we make a slightly larger incision through the back to directly remove bone spurs and soft tissue which are compressing your nerves prior to placing the screws and rods.

WHAT IS THE LIKELIHOOD THIS PROCEDURE WILL HELP MY PROBLEM?

This depends on your symptoms. The more buttock/leg/foot discomfort you have the more recovery you will have in general. Back pain is something that is difficult to predict will improve following surgery because there are many different causes of back pain that are not treated effectively with surgery. The goal of surgery is to take the pressure of the nerves. This helps very much with buttock and lower extremity symptoms but because there are many other causes of back pain that have nothing to do with the degenerative disks or pressure on the nerves, back pain relief is unpredictable following surgery. I generally advise patients that their buttock/lower extremity discomfort will improve significantly but any improvements in back pain is a bonus. The more lower extremity/buttock pain you have relative to back pain, the better patients tend to do. Patients will tend to have some aspect of back pain following this procedure since we must enter through the muscles of the back.

WHAT ARE THE RISKS OF SURGERY?

There are risks to any surgery. Life altering complications such as death, paralysis, and stroke are exceedingly rare. People generally have some degree of back pain following surgery due to stretch of the ligaments from creating more space between your bones with the cage or spacer. This varies by person. Because we are accessing your spine from the front or side, there is risk to the major blood vessels including the aorta and the vena cava. Injury to these blood vessels are exceedingly rare but can cause major complications including death. It is not uncommon to have some left upper thigh numbness that is usually temporary due to stretching of superficial sensory nerves during the surgery. There is a small risk for injury to the intestines as well. The risk of infection is less than 1% and typically occurs within the first couple weeks following surgery if it occurs. The odds of an infection rise with obesity, smoking, and poorly controlled diabetes. The goal of surgery is to fuse bones together. This generally takes about a year to complete. Fusion rate is upwards of 90%+ but patient

specific factors such as obesity, smoking, diabetes, and severe osteoporosis decrease those odds and may necessitate further surgery. The risk of future surgery at an area above or below the fusion is about 20%.

HOW LONG DOES THE SURGERY TAKE?

Surgery generally takes the morning once the procedure starts but it can vary when you add in the time for anesthesia to put you to sleep and wake you back up. Most of the time is positioning and set up and not total operative time.

WHAT IS THE FOLLOW-UP PROTOCOL?

We will have your follow-up at 6 weeks, 3 months, 7 months, and 1 year following surgery. If problems arise and you would like to be seen sooner, please contact our office. Occasionally we will close your wound with staples and if that occurs, we will see you back in the office at 2 weeks for staple removal.

HOW LONG IS THE RECOVERY AFTER SURGERY?

The overall recovery following surgery is about 1 year. The reason it is 1 year is that is about how long it takes for nerves to heal themselves to their full potential. Some people have permanent nerve damage from the degree or duration of compression that they have had and that may be a reason why full recovery is not achieved. But in general, how you feel at 1 year is how you will feel long-term. You will feel the initial post-operative soreness for at least a few weeks after surgery and that is generally how long it can take to return to desk/office type work. More physical work may require up to 6 weeks recovery to feel up to that level of activity. We will follow the progress of your fusion with x-rays, which takes about a year to complete.

WHAT ARE MY RESTRICTIONS AFTER SURGERY?

After surgery, it is generally no lifting more than 20 lbs until your first post-operative visit. This allows you to perform almost all usual household and self-care duties. The first 6 weeks are to allow your body to heal and recover from surgery so lifting more than that is not necessary, particularly bending and twisting.

DO I NEED A BRACE?

In most cases no. We will specifically speak to you if you do require a brace and give specific instructions on brace wear/care.

HOW MUCH PAIN WILL I HAVE AFTER SURGERY?

Your back/abdomen is going to hurt after surgery for a minimum of 4 weeks. This is from the muscle dissection required to take the pressure off your nerves. We will supply you with pain medication following surgery. It is not uncommon to have good days and bad days with your leg(s) following surgery. Some days you may feel great, other days you may feel like you did before surgery. This is part of the normal recovery process of the nerves. In general, pain improves first but other symptoms like numbness or weakness can take much longer to improve if at all.

WHEN CAN I DRIVE?

When you are off prescription pain medication/muscle relaxers during the day.

HOW DO I CARE FOR MY SURGICAL INCISION?

When you are leave the hospital, you will have already had your bandage changed. Please keep the bandage that you left the hospital with on for 3 days and then you can remove the bandage and shower normally. You will have butterfly strips over their incision which was sealed with glue. These strips will fall off as you shower. If they are not off in 1 week after surgery, please pull them off. Once you have showered for the first time, you should keep the incision covered for a week or two as it heals so it does not rub on your clothing especially if you have staples instead of butterfly strips. If you experience any drainage from the wound, please take a picture of the wound and email/call our office to evaluate it. If you have staples you can also shower 3 days after surgery. No baths or pools until you come into clinic for evaluation.

DO I NEED PHYSICAL THERAPY?

For the first 6 weeks after surgery, we want your body recover from surgery. Stick to the lifting restriction and refrain from frequent bending/twisting at the waist. If you would like to be involved in some gentle exercise such as going for a walk that is fine and you can walk as much as you feel comfortable. Formal physical therapy can be discussed at the first post-operative visit, but we need your muscles to heal.

WHAT WILL I NEED FOR PAIN MEDICATION?

Every patient should take 1000mg of Tylenol (2 extra strength over the counter Tylenols) 3x a day. This will help with giving you a background level of pain relief to decrease the amount of prescription pain medication you may need. We will also send you home with opioid pain medication and a muscle relaxer that you should take sparingly and only as needed (2-3x/a day maximum). If you are taking the opioids be sure to take a stool softener to prevent constipation. You may only need the opioids for a couple weeks. Do **not** take NSAIDs for 6 months following surgery since it decreases fusion rate.